CENTER FOR BEAM PHYSICS SEMINAR

"Synchronization by Passage Through Resonances in Non-Neutral Plasmas"

Lazar Friedland Hebrew University of Jerusalem

Friday, April 18, 2003, 10:30 AM Albert Ghiorso Conference Room (71-264), LBNL ••• Refreshments served at 10:20 AM •••

Abstract: Magnetized, pure-electron plasmas in a Malmberg-Penning trap can be modeled by two-dimensional Euler's equations of ideal fluids. The plasma density and electric potential in this approximation are analogous to the vorticity and stream function of the fluid velocity field. For example, a uniform-density electron plasma cylinder in a guiding magnetic field is equivalent to a circular vortex patch solution of an ideal fluid. Other, uniform, shape preserving m-fold symmetric plasma/vortex equilibria (V-states) where discovered by Deem and Zabusky. Typically, creation of these states requires some very nontrivial initial conditions on the plasma boundary. I will describe a different, more realizable (in experiments and simulations) approach to creation of V-states. The idea is similar to that used in controlling nonlinear waves in other extended systems and can be called "Pattern Formation by Synchronization". In the present application, we start from a trivial (axisymmetric) uniform plasma equilibrium, but consider a perturbed problem, where the perturbation is an oscillatory driving potential of appropriate spatial symmetry. We chirp the frequency of oscillations of the driving perturbation and pass through a resonance with the initial plasma equilibrium. Then, under certain conditions, the driven system is captured into a persistent nonlinear resonance, yielding a new, nontrivial rotating plasma state. I will also describe emergence and stability of nonuniform V-states.

<u>Biographical data and research interests</u>: Lazar Friedland is currently a Professor at the Hebrew University of Jerusalem on sabbatical leave at UC Berkeley. His research interests include basic plasma theory, linear and nonlinear waves in spacetime varying media, applications to resonances and mode conversion, pattern formation by adiabatic synchronization in dynamical and extended systems, non-equilibrium electron kinetics in plasmas, interaction of radiation and relativistic e-beams, and free electron lasers and accelerators.